

$C(O)OCH_2OC(O)R^{20}$, and $OCON(R^{20})_2$, $-CONR^7R^8$, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl, and heteroaryl, wherein the alkyl, alkenyl, alkynyl, aryl, heterocyclyl and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, NO_2 , heterocyclyl, aryl, heteroaryl, CF_3 , CN, OR^{20} , SR^{20} , $N(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $SO_2NR^{20}COR^{22}$, $SO_2NR^{20}CO_2R^{22}$, $SO_2NR^{20}CON(R^{20})_2$, $N(R^{20})_2$, $NR^{20}COR^{22}$, $NR^{20}CO_2R^{22}$, $NR^{20}CON(R^{20})_2$, $NR^{20}C(NR^{20})NHR^{23}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$, $CONR^{20}SO_2R^{22}$, $NR^{20}SO_2R^{22}$, $SO_2NR^{20}CO_2R^{22}$, $OCONR^{20}SO_2R^{22}$, $OC(O)R^{20}$, $C(O)OCH_2OC(O)R^{20}$, and $OCON(R^{20})_2$ and wherein the optional heteroaryl, aryl, and heterocyclyl substituent is optionally substituted with one or more substituents selected from the group consisting of halo, NO_2 , alkyl, CF_3 , amino, monoalkylamino, dialkylamino, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, $NCOR^{22}$, $NR^{20}SO_2R^{22}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$, $NR^{20}CON(R^{20})_2$, $OC(O)R^{20}$, $OC(O)N(R^{20})_2$, SR^{20} , $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, CN, and OR^{20} ;

R^5 and R^6 are each individually selected from the group consisting of H, and C_1 - C_{15} alkyl optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, NO_2 , heterocyclyl, aryl, heteroaryl, CF_3 , CN, OR^{20} , SR^{20} , $N(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $SO_2NR^{20}COR^{22}$, $SO_2NR^{20}CO_2R^{22}$, $SO_2NR^{20}CON(R^{20})_2$, $N(R^{20})_2$, $NR^{20}COR^{22}$, $NR^{20}CO_2R^{22}$, $NR^{20}CON(R^{20})_2$, $NR^{20}C(NR^{20})NHR^{23}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$, $CONR^{20}SO_2R^{22}$, $NR^{20}SO_2R^{22}$, $SO_2NR^{20}CO_2R^{22}$, $OCONR^{20}SO_2R^{22}$, $OC(O)R^{20}$, $C(O)OCH_2OC(O)R^{20}$, and $OCON(R^{20})_2$ and wherein the optional heteroaryl, aryl, and heterocyclyl substituent are optionally substituted with one or more substituents selected from the group consisting of halo, NO_2 , alkyl, CF_3 , amino, monoalkylamino, dialkylamino, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, $NCOR^{22}$, $NR^{20}SO_2R^{22}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$,

3
 cont'd

$\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{OC}(\text{O})\text{R}^{20}$, $\text{OC}(\text{O})\text{N}(\text{R}^{20})_2$, SR^{20} , $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, CN , and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl and heteroaryl, wherein the alkyl, alkenyl, alkynyl, aryl, heterocyclyl and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, NO_2 , heterocyclyl, aryl, heteroaryl, CF_3 , CN , OR^{20} , SR^{20} , $\text{N}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{SO}_2\text{NR}^{20}\text{COR}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{N}(\text{R}^{20})_2$, $\text{NR}^{20}\text{COR}^{22}$, $\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{C}(\text{NR}^{20})\text{NHR}^{23}$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{CONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{NR}^{20}\text{SO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{OCONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{OC}(\text{O})\text{R}^{20}$, $\text{C}(\text{O})\text{OCH}_2\text{OC}(\text{O})\text{R}^{20}$ and $\text{OCON}(\text{R}^{20})_2$ and wherein the optional heteroaryl, aryl and heterocyclyl substituents are optionally substituted with one or more substituents selected from the group consisting of halo, NO_2 , alkyl, CF_3 , amino, monoalkylamino, dialkylamino, llylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, NCOR^{22} , $\text{NR}^{20}\text{SO}_2\text{R}^{22}$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{OC}(\text{O})\text{R}^{20}$, $\text{OC}(\text{O})\text{N}(\text{R}^{20})_2$, SR^{20} , $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, CN , and OR^{20} ;

R^8 is selected from the group consisting of hydrogen, C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl, and heteroaryl, wherein the alkyl, alkenyl, alkynyl, aryl, heterocyclyl, and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, NO_2 , heterocyclyl, aryl, heteroaryl, CF_3 , CN , OR^{20} , SR^{20} , $\text{N}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{SO}_2\text{NR}^{20}\text{COR}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{N}(\text{R}^{20})_2$, $\text{NR}^{20}\text{COR}^{22}$, $\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{C}(\text{NR}^{20})\text{NHR}^{23}$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{CONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{NR}^{20}\text{SO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{OCONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{OC}(\text{O})\text{R}^{20}$, $\text{C}(\text{O})\text{OCH}_2\text{OC}(\text{O})\text{R}^{20}$, and $\text{OCON}(\text{R}^{20})_2$ and wherein each optional heteroaryl, aryl, and heterocyclyl substituent is optionally substituted with one or more substituents selected from

the group consisting of halo, NO₂, alkyl, CF₃, amino, monoalkylamino, dialkylamino, lkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, NCOR²², NR²⁰SO₂R²², COR²⁰, CO₂R²⁰, CON(R²⁰)₂, NR²⁰CON(R²⁰)₂, OC(O)R²⁰, OC(O)N(R²⁰)₂, SR²⁰, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, CN, and OR²⁰;

R²⁰ is selected from the group consisting of H, C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl, and heteroaryl, wherein the alkyl, alkenyl, alkynyl, heterocyclyl, aryl, and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, monoalkylamino, dialkylamino, lkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, CN, O-C₁₋₆ alkyl, CF₃, aryl, and heteroaryl;

*a3
cont'd*
R²² is selected from the group consisting of C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl, and heteroaryl, wherein the alkyl, alkenyl, alkynyl, heterocyclyl, aryl, and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, monoalkylamino, dialkylamino, lkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, CN, O-C₁₋₆ alkyl, CF₃, aryl, and heteroaryl; and

wherein R² and R⁴ are selected from the group consisting of H, C₁₋₆ alkyl, and aryl that is optionally substituted with halo, CN, CF₃, OR²⁰ and N(R²⁰)₂, with the proviso that when R² is not hydrogen then R⁴ is hydrogen, and when R⁴ is not hydrogen then R² is hydrogen.

2. (Once Amended) The compound of claim 1 wherein R³ is selected from the group consisting of C₁₋₁₅ alkyl, halo, CF₃, CN, OR²⁰, SR²⁰, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, COR²⁰, CO₂R²⁰, -CONR⁷R⁸, aryl and heteroaryl wherein the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group

consisting of halo, aryl, heteroaryl, CF_3 , CN , OR^{20} , SR^{20} , S(O)R^{22} , SO_2R^{22} , $\text{SO}_2\text{N(R}^{20})_2$, COR^{20} , CO_2R^{20} and $\text{CON(R}^{20})_2$, and wherein each optional heteroaryl and aryl substituent is optionally substituted with one or more substituents selected from the group consisting of halo, alkyl, CF_3 , CN , and OR^{20} ;

R^5 and R^6 are each individually selected from the group consisting of H, and $\text{C}_1\text{-C}_{15}$ alkyl optionally substituted with one aryl substituent that is optionally substituted with one or more substituents selected from the group consisting of halo and CF_3 ;

R^7 is selected from the group consisting of C_{1-15} alkyl, C_{2-15} alkynyl, aryl, and heteroaryl, wherein the alkyl, alkynyl, aryl, and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, aryl, heteroaryl, CF_3 , CN , and OR^{20} , and wherein each optional heteroaryl and aryl substituent is optionally substituted with one or more substituents selected from the group consisting of halo, alkyl, CF_3 , CN , and OR^{20} ;

R^8 is selected from the group consisting of hydrogen and C_{1-15} alkyl;

R^{20} is selected from the group consisting of H, C_{1-4} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with one alkyl substituent; and

R^{22} is selected from the group consisting of C_{1-4} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with from 1 to 3 alkyl groups.

3. (Once Amended) The compound of claim 1 wherein R^3 is selected from the group consisting of C_{1-15} alkyl, halo, CF_3 , CN , OR^{20} , CO_2R^{20} , $-\text{CONR}^7\text{R}^8$, aryl and heteroaryl, wherein the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, aryl, CF_3 , CN , OR^{20} , CO_2R^{20} and $\text{CON(R}^{20})_2$, and wherein each optional heteroaryl and aryl substituent is optionally substituted

a3
cont'd

X

with one or more substituents selected from the group consisting of halo, alkyl, CF₃, CN, and OR²⁰;

R⁵ and R⁶ are each individually selected from the group consisting of hydrogen and C₁₋₆ alkyl;

R⁷ is selected from the group consisting of C₁₋₁₀ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, heteroaryl, CF₃, CN, and OR²⁰, and wherein each optional heteroaryl and aryl substituent is optionally substituted with at least one substituent selected from the group consisting of halo, alkyl, CF₃, CN, and OR²⁰;

R⁸ is selected from the group consisting of hydrogen and C₁₋₁₅ alkyl; and

R²⁰ is selected from the group consisting of hydrogen and C₁₋₄ alkyl.

3
a
cont'd

4. (Once Amended) The compound of claim 1 wherein R³ is selected from the group consisting of C₁₋₁₀, alkyl, halo, CF₃, CN, CO₂R²⁰, -CONR⁷R⁸, aryl and heteroaryl wherein the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, CF₃, CN, OR²⁰ and CON(R²⁰)₂;

R⁵ and R⁶ are each individually selected from the group consisting of hydrogen and C₁₋₆ alkyl;

R⁷ is selected from the group consisting of C₁₋₁₀ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, heteroaryl, CF₃, CN, OR²⁰ and wherein each optional heteroaryl and aryl substituent is optionally substituted with at least one substituent selected from the group consisting of halo, alkyl, CF₃, CN, and OR²⁰;

R⁸ is selected from the group consisting of hydrogen and C₁₋₁₅ alkyl; and

A

R^{20} is selected from the group consisting of hydrogen and C_{1-4} alkyl.

5. (Once amended) The compound of claim 1 wherein R^3 is selected from the group consisting of C_{1-10} alkyl, halo, CF_3 , CN, OR^{20} , CO_2R^{20} , $-CONR^7R^8$ and aryl wherein the alkyl and aryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, CF_3 , CN, OR^{20} and $CON(R^{20})_2$;

R^5 and R^6 are each individually selected from the group consisting of hydrogen and C_{1-6} ;

R^7 is selected from the group consisting of C_{1-10} alkyl, aryl, and heteroaryl, where the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, heteroaryl, CF_3 , CN, and OR^{20} and wherein each optional heteroaryl and aryl substituent is optionally substituted with at least one substituent selected from the group consisting of halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from the group consisting of hydrogen and C_{1-15} alkyl; and

R^{20} is selected from the group consisting of hydrogen and C_{1-4} alkyl.

6. (Once amended) The compound of claim 1 wherein R^1 is $-CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$ and aryl; wherein the aryl substituent is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, C_{1-6} alkyl, CF_3 , CN, OR^{20} , and $CON(R^{20})_2$;

R^7 is selected from the group consisting of hydrogen, C_{1-10} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, CF_3 , CN, and OR^{20} and wherein each optional aryl substituent is optionally substituted with at least one substituent selected from the group consisting of halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from the group consisting of hydrogen and C_{1-15} alkyl; and

3
contd

X

R^{20} is selected from the group consisting of hydrogen and C_{1-4} alkyl.

7. (Once amended) The compound of claim 1 wherein R^1 is $-CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$ and aryl wherein the aryl substituent is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, C_{1-6} alkyl, CF_3 and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, C_{1-8} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with one substituent selected from the group consisting of halo, aryl, CF_3 , CN, and OR^{20} and wherein each optional aryl substituent is optionally substituted with at least one substituent selected from the group consisting of halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from the group consisting of hydrogen and C_{1-8} alkyl; and

R^{20} is selected from the group consisting of hydrogen and C_{1-4} alkyl.

8. (Once amended) The compound of claim 1 wherein R^1 is $-CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$, and aryl that is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, C_{1-3} alkyl, CF_3 and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, and C_{1-8} alkyl that is optionally substituted with one substituent selected from the group consisting of halo, CF_3 , CN and OR^{20} ;

R^8 is selected from the group consisting of hydrogen and C_{1-3} alkyl; and

R^{20} is selected from the group consisting of hydrogen and C_{1-4} alkyl.

9. (Once amended) The compound of claim 1 wherein R^1 is $-CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$, and aryl that is optionally substituted with one substituent selected from the group consisting of halo, C_{1-3} alkyl, and OR^{20} ;

X

Sub B
 R^7 is selected from the group consisting of hydrogen, and C_{1-3} alkyl that is optionally substituted with one substituent selected from the group consisting of halo, CF_3 , CN and OR^{20} ;

R^8 is hydrogen; and

R^{20} is selected from the group consisting of hydrogen and C_{1-4} alkyl.

10. (Once amended) The compound of claim 1 wherein R^1 is $-CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$, and aryl that is optionally substituted with one substituent selected from the group consisting of halo, C_{1-3} alkyl and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, and C_{1-3} alkyl;

R^8 is hydrogen; and

R^{20} is selected from the group consisting of hydrogen and C_{1-4} alkyl.

5 11. (Once amended) The compound of claim *10* wherein R^7 is a methyl.

6 12. (Once amended) The compound of claim *10* wherein R^3 is $-CO_2Et$.

13. (Once amended) The compound of claim 1 wherein R^1 is $-CONHET$;

3
 R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$, and aryl that is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, C_{1-6} alkyl, CF_3 , CN, OR^{20} , and $CON(R^{20})_2$;

R^7 is selected from the group consisting of hydrogen, C_{1-10} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, CF_3 , CN, and OR^{20} and wherein each optional aryl substituent is optionally substituted with at least one substituent selected from the group consisting of halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from the group consisting of hydrogen, and C_{1-15} alkyl; and

R^{20} is selected from the group consisting of hydrogen, and C_{1-4} alkyl.

14. (Once amended) The compound of claim 1 wherein R^1 is $-\text{CONH}Et$;

R^3 is selected from the group consisting of CO_2R^{20} , $-\text{CONR}^7R^8$, and aryl that is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, C_{1-6} alkyl, CF_3 and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, C_{1-8} alkyl, and aryl, wherein the alkyl and aryl substituents are optionally substituted with one substituent selected from the group consisting of halo, aryl, CF_3 , CN, and OR^{20} and wherein each optional aryl substituent is optionally substituted with one or more substituents selected from the group consisting of halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from the group consisting of hydrogen, and C_{1-8} alkyl; and

R^{20} is selected from the group consisting of hydrogen, and C_{1-4} alkyl.

15. (Once amended) The compound of claim 1 wherein R^1 is $-\text{CONH}Et$;

R^3 is selected from the group consisting of CO_2R^{20} , $-\text{CONR}^7R^8$, and aryl that is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, C_{1-3} alkyl, CF_3 and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, and C_{1-8} alkyl that is optionally substituted with one substituent selected from the group consisting of halo, CF_3 , CN and OR^{20} ;

R^8 is selected from the group consisting of hydrogen, and C_{1-3} alkyl; and

R^{20} is selected from the group consisting of hydrogen, and C_{1-4} alkyl.

16. (Once amended) The compound of claim 1 wherein R^1 is $-\text{CONH}Et$;

R^3 is selected from the group consisting of CO_2R^{20} , $-\text{CONR}^7R^8$, and aryl that is optionally substituted with one substituent selected from the group consisting of halo, C_{1-3} alkyl and OR^{20} ;

3
contd

X

R⁷ is selected from the group consisting of hydrogen and C₁₋₃ alkyl that is optionally substituted with one substituent selected from halo, CF₃, CN and OR²⁰;

R⁸ is hydrogen; and

R²⁰ is selected from the group consisting of hydrogen, and C₁₋₄ alkyl.

17. (Once amended) The compound of claim 1 wherein R¹ is -CONHET;

R³ is selected from the group consisting of CO₂R²⁰, -CONR⁷R⁸, and aryl that is optionally substituted with one substituent selected from the group consisting of halo, C₁₋₃ alkyl and OR²⁰;

R⁷ is selected from the group consisting of hydrogen, and C₁₋₃ alkyl;

R⁸ is hydrogen; and

R²⁰ is selected from the group consisting of hydrogen, and C₁₋₄ alkyl.

18. (Once amended) The compound of claim 10 where R¹ is -CONHET.

19. (Once amended) A method for stimulating coronary vasodilation in a mammal by administering to the mammal a therapeutically effective amount of a compound of claim 1 that is sufficient to stress the heart and induce a coronary steal situation for the purposes of imaging the heart.

20. (Once amended) A pharmaceutical composition comprising the composition of claim 1 and one or more pharmaceutical excipients;

21. (Once amended) The pharmaceutical composition of claim 20 wherein the pharmaceutical composition is in the form of a solution.

Add the following new claims 26-28:

26. (New) The compound of claim 1 selected from the group consisting of N⁶-{3-(3R)tetrahydrofuranyl}-2-(N-1-(4-ethoxycarbonyl)pyrazolyl)adenosine; N⁶-{3-(3R)tetrahydrofuranyl}-2-(N-1-(4-(4-chlorophenyl))pyrazolyl)adenosine; N⁶-{3-

35